RESEARCH ARTICLE

Continuation and user satisfaction of the levonorgestrel intrauterine system (LNG IUS) contraceptive in Nigeria

[version 1; peer review: awaiting peer review]

Anthony Nwala , Ekerette Udoh , Jennifer Anyanti, Adewole Fajemisin

Department of Research, Family Planning and Reproductive Health, Society for Family Health, Abuja, Nigeria

V1 First published: 02 Feb 2022, 6:4
<https://doi.org/10.12688/gatesopenres.13195.1>
Latest published: 02 Feb 2022, 6:4
<https://doi.org/10.12688/gatesopenres.13195.1>

Open Peer Review

Approval Status Awaiting Peer Review

Any reports and responses or comments on the article can be found at the end of the article.

Abstract

Background: The hormonal intrauterine device or intrauterine system (IUS) also known as the levonorgestrel intrauterine system (LNG-IUS) is a highly effective hormonal intra-uterine contraceptive. However, services offering the method are not widely available in Nigeria and little evidence exists on the dynamics of its use. We examined the continuation rate and satisfaction with the IUS among the user population.

Methods: This prospective longitudinal phone survey involved a baseline survey of users at two-weeks post-insertion of the LNG IUS, recruited from 40 clinics across 17 states in Nigeria, with a follow-up survey at the 3rd and 12th months. A total of 208 users were interviewed at baseline, 98 at three months, and 73 at 12 months. User family planning and the IUS use experiences were elicited, as well as the continuation rate and satisfaction with the method at three and 12 months.

Results: At three- and 12 months post-insertion, 96.9% (95% CI: 91.3, 99.3) and 91.8% (95% CI: 82.9, 96.9), respectively, reported still using the LNG IUS, with none out of the few users who discontinued the method reporting a method failure. Discontinuation was mainly a result of the experience of menstrual bleeding or amenorrhea (25.0%), experiences of pain with the method (18.8%), and partner complaining about strings (16.7%). High satisfaction with the LNG IUS (76.5% at three months and 86.3% at 12 months post-insertion) was reported. Satisfaction with LNG IUS was significantly associated with not having breast tenderness/pain (88.2%) and no vaginal bacterial infection (87.5%) at 12 months compared to experiencing breast tender/pain (50.0%) and vaginal infection (0.0%) ($p < 0.05$).

Conclusion: High user continuation and satisfaction with IUS indicates the positive potential of the method as a contraceptive in Nigeria.

Keywords

Levonorgestrel Intrauterine System (LNG IUS), contraceptive method discontinuation, contraceptive method mix, Nigeria Family Planning.

Corresponding author: Ekerette Udoh (eudoh@sfnigeria.org)

Author roles: **Nwala A:** Conceptualization, Investigation, Methodology, Project Administration, Resources, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Udoh E:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Anyanti J:** Funding Acquisition, Methodology, Project Administration, Resources, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Fajemisin A:** Investigation, Methodology, Project Administration, Resources, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This study was made possible by the support of the American People through the United States Agency for International Development (USAID). The SIFPO2 project [cooperative agreement number AID-OAA-A-14-00037] implemented this research in Nigeria. SFH also has an affiliation with Bill and Melinda Gates Foundation implementing FP services expansion project in the grant: Integrating FP & PHC Services Expansion in Community Pharmacies & Patent Proprietary Medicine Vendors [OPP1173898]. This article publication fee was supported through this grant by Bill and Melinda Gates Foundation.

Copyright: © 2022 Nwala A *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Nwala A, Udoh E, Anyanti J and Fajemisin A. **Continuation and user satisfaction of the levonorgestrel intrauterine system (LNG IUS) contraceptive in Nigeria [version 1; peer review: awaiting peer review]** Gates Open Research 2022, 6:4 <https://doi.org/10.12688/gatesopenres.13195.1>

First published: 02 Feb 2022, 6:4 <https://doi.org/10.12688/gatesopenres.13195.1>

Introduction

Discontinuation of contraception among women in Nigeria is one of the main challenges with contraceptive use, together with generally low use of contraception. The recent national demographic health survey report (NDHS 2018) indicates that about 41% of women who had an episode of use of contraception discontinued the contraception within 12 months of use¹. Contraceptive discontinuation and dissatisfaction with a method occur due to several reasons which may be method related. While the desire to become pregnant is one of the major reasons why a woman will discontinue contraception, method-related reasons such as side effects, method failure, and the need for a more effective method are also prominent reasons for method discontinuation among women who still wish to prevent pregnancy. Discontinuation of contraceptives often leads to unwanted pregnancies and reduces the effort of family planning (FP) programs². Studies show that a sizable proportion of women become exposed to the risk of unintended births after discontinuation of contraceptives^{3,4}. In Nigeria, it has been reported that an estimated 15.6% of unintended pregnancies are among women who discontinued contraception in the last seven months⁴. Even higher proportions of unintended births have also been reported in other countries as a result of the discontinuation of contraceptives in the last three months³. Serious reproductive consequences of discontinuation because of reported method failure and method-related reasons can occur. Accidental pregnancies that end in miscarriage, stillbirth, or abortion have been reported³.

In many developing countries including Nigeria, many health outlets providing services for modern contraception are highly constrained on the method mix available for potential users⁵. It is widely known that limited method choice can hinder the uptake and use of contraception, and has been linked with dissatisfaction and discontinuation of contraception^{6,7}. The modern contraceptive method mix in Nigeria comprises mainly injectable contraceptives (33.8%), pills (23%), male condoms (18.1%), Implant (10.9%), and intra-uterine devices (IUDs) (5.2%). Other methods such as the lactational amenorrhea method (LAM) make up about 4.8%, while female and male sterilization make up 1.6% and 0.2% of the modern contraceptive method mix, respectively⁸. Global data is available on the dynamics of contraceptive use as it relates to the discontinuation of methods. According to data from 19 different Demographic Health Survey (DHS) participating countries, an average of 38% of women reported discontinuing their use of reversible methods by the 12th month and 64% by the 36th month³. The lowest 12-month discontinuation rate was noted for the intrauterine device (IUD; 13%) and the highest was for condoms (50%), while the pill and injectable contraceptives were discontinued by about 40% of users within the first 12 months of use. Contraceptive continuation indicates the acceptability of a contraceptive method, and contraceptive continuation rate is the cumulative probability that acceptors of a contraceptive method will still be using any contraceptive method offered by a program after a specified period⁹.

Given this understanding, a high continuation rate of a method indicates users' high level of compliance with the method.

From the pattern of contraceptive method-use according to the Nigeria 2018 National Demographic Health Survey, the national prevalence of 5% of IUD users, use mainly the copper IUD, which is the most common intrauterine contraceptive available in FP clinics in Nigeria. Meanwhile, other types of intrauterine devices such as the hormonal IUD which is a progestin-based intra-uterine delivery system and is a long-acting and reversible contraceptive (LARC), also exist. However, the hormonal IUD does not contribute to the proportion of intrauterine contraceptives used by women¹⁰. In one study in Zaria Nigeria, of all the 1104 women who opted for an IUD as a method of contraception only 68 (6.1%) women chose the hormonal IUD¹¹. This is in contrast to other countries; for example in the US, where the proportion of IUD users has been reported as 12%, with 74% of the users of the intrauterine contraceptives choosing the hormonal IUD¹². In Nigeria, the use of the hormonal IUD is limited and is widely unavailable to users of contraceptives from both public and private clinics. The limited use of the method is mainly due to its high cost; a unit of the LNG IUS brand Mirena®, which is manufactured by Bayer Healthcare Pharmaceuticals Inc., costs between 25,000 to 90,000 Nigerian Naira, which is currently equivalent to about \$69 – \$250, compared with the copper IUD, which costs about \$5 or less at private clinics and is free of charge in government clinics. In addition, the hormonal IUD method has been relatively recently introduced in the country despite its decades of existence since its development in the 1970s^{4,13}. There is also the issue of the availability of providers who are trained to provide the method, thus limiting its use.

The hormonal IUD is a T-shaped device comprising a cylinder containing 52 mg of levonorgestrel, with an average release of 14µg of levonorgestrel every 24 hours of the life of the IUS¹⁴. The Mirena® LNG IUS that is produced by Bayer Healthcare Pharmaceuticals Inc is indicated to have an effective duration of up to 5 years. However, clinical trials suggest that the LNG IUS can provide contraceptive protection for up to 7 years^{15,16}. From a recent study trial, a brand of the IUS Lilelleta developed by Medicine360 (registered under the trade name as Avibela™ in the FP2020 countries including Nigeria), which was previously labeled for use for 3 years has also been indicated to have an effective duration over five years¹⁷. Eliora another brand of the hormonal IUD developed by Pregna is similar to the Mirena having an effective duration of 5 years¹⁸. The IUS functions locally in the uterus by thickening the cervical mucus and suppressing endometrial proliferation, thus inhibiting conception. The IUS is known to have both contraceptive and non-contraceptive health benefits in the treatment of menorrhagia, endometriosis, and endometrial hyperplasia¹⁴. The method is very convenient as it requires nothing else to be done regularly to maintain contraception until after five years when the device can be replaced.

Most side effects of the IUS are transient and include changes in menstrual bleeding pattern, headache, abdominal cramps, breast tenderness, and weight gain¹³.

To expand the range of contraceptive options available in the country and to improve the acceptability of contraception, the introduction of new and effective methods such as the hormonal IUS as a strategy to increase contraceptive uptake has been the focus of some FP interventions and organizations that are piloting roll out of new and affordable IUS to potential users in Nigeria. For instance, since 2017 the Society for Family Health (SFH) has been supporting service provision for the LNG IUS as part of a broad range of options available in 40 private health clinics across 17 states in Nigeria, using donations of the registered product in Nigeria from the International Contraceptive Access (ICA) Foundation. To better understand the potential of the hormonal IUS method to contribute to the contraceptive method mix in Nigeria, this study was therefore undertaken to assess and document the continuation of use of the LNG IUS and satisfaction with the method, as well as user characteristics and experiences that may influence continuation and satisfaction among the user population. This will add to the existing body of knowledge on the method and document the experiences of users of the method in Nigeria. If users show high satisfaction and continuation with the IUS, these findings will go a long way to encourage adoption and utilization of the method by users and provide the motivation for its recommendation by FP providers and clinicians, both as a contraceptive and for its non-contraceptive benefits such as treatment of menorrhagia and other gynecological disorders.

Methods

Ethical statement

The study protocol was reviewed by the National Health Research Ethics committee in 2017 and ethical approval was obtained (NHREC/01/01/2007-05/02/2017). Given that data collection in this study was done via phone interviews, oral informed consent was deemed adequate and was obtained from all participants. The consent to partake in the study was recorded by the researchers along with responses of respondents and a written consent form was signed by the interviewers for documentation.

Study design and setting

This study was a prospective longitudinal survey of 209 women of reproductive age 18 – 49 years who received the LNG IUS as a method of choice after undergoing counseling using the Balanced Counselling Strategy (BCS) for FP. The study was conducted in 40 select private health clinics that are members of the SFH Healthy Family Network (HFN), a social franchise of private health facilities. The 40 selected facilities had high numbers of clientele with IUDs and are implementing service delivery for LNG IUS under the Supporting International Family Planning Organisation (SIFPO) Project, which provided the IUS method in the context of informed choice using the ICA Foundation donated LNG IUS product. The facilities are in 17 states in Nigeria, namely Abia, Akwa

Ibom, Cross River, Enugu, Rivers, Kano, Katsina, Imo, Gombe, Taraba, Lagos, Ogun, Oyo, Benue, Niger, Edo, and the Federal Capital Territory (FCT) where the franchise operates.

Participant selection

Trained health providers in the facilities recruited consenting women who already accepted the IUS as their method of choice to participate in a telephone interview. Only women aged 18 to 49 years were recruited. Recruited participants were contacted by trained call agents/interviewers at predetermined periods: at two weeks post-insertion of the IUS in the facility, at three months, and after 12 months. Women who consented and agreed for their phone number to be passed on to the research team were contacted for interviews and follow-up. Only their first names and phone numbers were passed to the research team via SMS. The SMS was immediately deleted from the provider's phones after receipt from the research team. Women less than 18 years of age were excluded from this study. Confidentiality of the respondent's information was ensured by assigning unique identification codes to each recruited participant, and all respondents' data were electronically stored, and the files were password protected. The sample size calculation for the study was based on the following formula:

$$N = \frac{[Z_{1-\alpha}\sqrt{2p(100-p)} + Z_{1-\beta}\sqrt{p_1(100-p_1) + p_2(100-p_2)}]^2}{(p_2 - p_1)^2}$$

Where N is the sample size, P1 (proportion of the sample reporting satisfaction at time 1) = 75%, P2 (proportion of the sample reporting satisfaction at time 2) = 60%, Z1- α is the standard normal deviate corresponding to a two-sided level of significance (α) of 5%, Z1-β (statistical power) = 0.8. The initial calculated sample size was 135 considering a design effect of 1.2. The final sample of 208 was recruited with the consideration of about 45% loss-to-follow-up rate between baseline and the 12-month follow-up.

Data collection

Women were considered as loss-to-follow up if they could not be reached after at least five repeated attempts within a couple of days. The loss-to-follow-up was mainly due to non-reachable phone contacts after five repeated attempts and due to women refusing interviews during the follow-up. No form of incentive was given to participants in the study. Baseline data collection started in May 2017 and lasted for about 8 months until the required sample was achieved. Subsequent follow-up interviews were done on a date scheduled with the participant at three months and 12 months. Data collection, therefore, ended in 2018 after the last scheduled 12-month interview. Data were collected electronically using a questionnaire implemented on the ODK android software. The ODK is a computer-assisted personal interview (CAPI) application available for Android devices for electronic data collection.

Information was collected on the respondent's background characteristics and their FP use experience, which included

pregnancy intention, the occasion for the insertion of the IUS such as miscarriage or abortion, side effects (experienced during the use of IUS at three months and 12 months) and complaints of discomfort from the IUS string by her partner. The women were also asked about how satisfied they were with using the IUS, and if they would recommend the IUS to other women or a friend. A Likert scale was used to score the responses of the women to the satisfaction question. The Likert scale ranged from 1–5, with an increasing score indicating increasing satisfaction with the method. User satisfaction response was re-coded such that a respondent's score of 1–3 indicated no satisfaction, while scores of 4–5 indicated satisfaction with the method. Continuation rate was measured as the proportion of women who received the IUS and reported they are still using the method at the time of the three-month and 12-month follow-up surveys.

Statistical analysis

Data was presented in this study using descriptive statistics with frequencies, percentages, and 95% confidence intervals as appropriate. Association between independent variables of background characteristics and FP/ IUS experiences with the dependent variables (continuation of the IUS and satisfaction) were tested using the Chi-square test. Statistical significance for association was set at $p < 0.05$. Given the attrition recorded in the survey, possible attrition bias was examined by carrying out a Chi-square test to compare the background characteristics between those who were reached in the second and third rounds of the study with those who were not reached. Statistically significant differences in the background characteristics between those reached and those not reached would indicate some existence of attrition bias in the result estimates. Data analysis was done using IBM SPSS Statistics - version 25.

Results

User background characteristics, FP and IUS use experiences

A total of 208 users were interviewed at baseline. At the three-month follow-up, 98 women were successfully reached and interviewed, while at the 12-month follow-up, 73 women were successfully reached and interviewed. The average age of users of the IUS in the study at baseline was 33.7 ± 6 , and the majority age group was 35 years and older (44.2%), while approximately 30% and 20% of the users were aged between 30–34 years and 25–29 years old, respectively. More than half of users had between 1–3 children (58.2%), another good proportion, 38.5%, had 4–7 children, and very few users had no children (1.4%) (Table 1). The Chi-square test results in Table 1 showed statistical significance for only age at 3-month for the sample due to the attrition. For all other background characteristics, no statistical significance was observed in the background characteristics of the participants reached and those not reached in the follow-up surveys. Nearly a third of the women in the study reported they were no longer seeking pregnancy (31.7%), while another 23.6% said they would seek a pregnancy in the next 3–5 years, and 10% said in the next 1–2 years.

Around 4% reported receiving the IUS two weeks before carrying out an abortion or after a miscarriage. Up to 96.2% of users of the method reported they were satisfied with getting the IUS inserted at the clinic at the uptake of the method. Table 2 presents more of the IUS use experiences and characteristics of the participants.

The most common side effects were less or no bleeding, irregular bleeding, and breast tenderness/pain. Reports of less bleeding increased from 8.2% at three months to 24.7% at 12 months, while irregular bleeding which was high at 3 months (32.7%) reduced to 4.1% at 12 months. The prevalence of other side effects during the use of the IUS at three months and 12 months are presented in Table 3.

Some of the reasons for discontinuation of the IUS at 12 months were mainly due to experience of decreased menstrual bleeding or amenorrhea (25%), pain (18%), and weight gain (12%). Other reasons were as a result of worry of being pregnant because of no bleeding (6.3%) and partner complaining about feeling the strings of the IUS (16.7) (see Figure 1).

Concerning recommending the benefits of the IUS, a high rate for a recommendation of the benefits of the IUS were seen for attributes such as its fewer side effects (23.4% at 3 months and 14.4% at 12 months) and its convenience of use (21.9% at 3 months and 15.1% at 12 months). Other reported high reasons for recommending the IUS were its high effectiveness, long duration of effectiveness, reduced bleeding, and affordability of the product at the facility (Table 4).

Rates of continuation and user satisfaction with the IUS

Of the 98 women who were interviewed at three months post-insertion of their IUS, 96.9% (95% CI: 91.3, 99.3) of them were still using the method, while at 12 months post-insertion, 91.8% (95% CI: 82.9, 96.9) of the 73 women reached reported still using the IUS method. The proportion of users who were satisfied with the method at baseline at two weeks post-uptake was 95.2% (95% CI: 91.3, 97.6), while satisfaction with the method at three months and 12 months post-uptake was 76.5% (95% CI: 66.8, 84.5) and 86.3% (95% CI: 77.5, 94.1), respectively.

Continuation and satisfaction with IUS according to user characteristics and experiences

Our analysis showed that continuation of IUS at 12 months was significantly lower (83.8%) among women who visited a provider because of problems with the method than women who did not visit a provider (100%). Analyses were also conducted to see if continuation was correlated with age, fertility intention, counseling, etc. but none of these were found to be statistically significant. In addition to the higher rate of uptake among older women, the continuation rate at 12 months was higher among older women of ages 30 years and older, ranging from about 92% to about 97%, compared with those aged 25 – 29 years old, which was about 80% (Table 5). According to marital status, high rates for the continuation of the IUS

Table 1. Baseline demographic characteristics of users of levonorgestrel intrauterine system and the assessment of attrition in the follow-up surveys.

Background characteristics	Survey at baseline	Survey at 3 months follow-up			Survey at 12 months follow-up		
		Dropped out	Participated	χ^2 (p)	Dropped out	Participated	χ^2 (p)
	n (%)	n (%)	n (%)		n (%)	n (%)	
Age category							
18–24 years old	12 (5.8)	11 (91.7)	1 (8.3)	7.906 (0.048)	11 (91.7)	1 (8.3)	5.975 (0.113)
25 – 29 years old	42 (20.2)	21 (50.0)	21 (50.0)		27 (64.3)	15 (35.7)	
30 – 34 years old	62 (29.8)	30 (48.4)	32 (51.6)		35 (56.5)	27 (43.5)	
35 and older	92 (44.2)	48 (52.2)	44 (47.8)		62 (67.4)	30 (32.6)	
Highest level of schooling							
Never attended school	2 (1.0)	1 (50.0)	1 (50.0)	5.099 (0.165)	2 (100.0)	0 (0.0)	3.783 (0.286)
Attended but did not complete school	4 (1.9)	4 (100.0)	0 (0.0)		4 (100.0)	0 (0.0)	
Primary	16 (7.7)	6 (37.5)	10 (62.5)		9 (56.2)	7 (43.8)	
Secondary or higher	186 (89.4)	99 (53.2)	87 (46.8)		120 (64.5)	66 (35.5)	
Marital status							
Single	3 (1.4)	3 (100.0)	0 (0.0)	3.836 (0.280)	3 (100.0)	0 (0.0)	3.929 (0.269)
Married/living together	198 (95.2)	102 (51.5)	96 (48.5)		126 (63.6)	72 (36.4)	
Widowed	3 (1.4)	2 (66.7)	1 (33.3)		2 (66.7)	1 (33.3)	
Divorced/separated	4 (1.9)	3 (75.0)	1 (25.0)		4 (100.0)	0 (0.0)	
Religion							
Islam	37 (17.8)	23 (62.2)	14 (37.8)	3.846 (0.146)	28 (75.7)	9 (24.3)	5.363 (0.068)
Christian (non-Catholic)	151 (72.6)	80 (53.0)	71 (47.0)		98 (64.9)	53 (35.1)	
Christian (Catholic)	20 (9.6)	7 (35.0)	13 (65.0)		9 (45.0)	11 (55.0)	
Parity							
0	3 (1.4)	3 (100.0)	0 (0.0)	5.133 (0.162)	3 (100.0)	0 (0.0)	2.098 (0.552)
1–3	121 (58.2)	60 (49.6)	61 (50.4)		76 (62.8)	45 (37.2)	
4–7	80 (38.5)	46 (57.5)	34 (42.5)		53 (66.2)	27 (33.8)	
8 and more	4 (1.9)	1 (25.0)	3 (75.0)		3 (75.0)	1 (25.0)	
Age of youngest child							
Less than 1 year old	75 (36.6)	40 (53.3)	35 (46.7)	4.138 (0.126)	47 (62.7)	28 (37.3)	3.416 (0.181)
1–5 years	104 (50.7)	49 (47.1)	55 (52.9)		64 (61.5)	40 (38.5)	
6 years and above	26 (12.7)	18 (69.2)	8 (30.8)		21 (80.8)	5 (19.2)	
Total	208 (100.0)	107 (52.2)	98 (47.8)		135 (64.9)	73 (35.1)	

were seen at three and 12 months post-insertion among women living with their partner, which was 96.8% at 3 months and

91.7% at 12 months. As there were not enough single women in the study, the continuation rate among this group was not

Table 2. Family planning and experiences of levonorgestrel intrauterine system (IUS) use.

	Frequency	Percent
Pregnancy intention (baseline)		
In less than a year	1	0.5
In 1–2 years	21	10.1
In 3–5 years	49	23.6
In more than 5 years	12	5.8
Never	66	31.7
Do not know	59	28.4
Receiving IUS within two weeks of having an abortion or miscarriage (baseline)		
Yes	8	3.8
No	200	96.2
Satisfaction in getting the IUS inserted (baseline)		
Yes	200	96.2
No	8	3.8
Impact of not bleeding to wellbeing at 12-month		
Positive	5	27.8
Negative	7	38.9
Neutral	6	33.3
Partner feels IUS strings at 12 months		
Yes	18	24.7
No	53	72.6
Not sure	2	2.7
Counselling at uptake about what to do if side-effect is experienced (baseline)		
Yes	68	32.5
No	135	64.9
Do not know	5	2.4
Visits to provider because of problem with method at 12 months		
Yes	37	50.7
No	36	49.3
Likelihood of recommending the IUS at 12 weeks post insertion		
Would recommend	200	96.2
Would not recommend	2	1.0
Not sure	6	2.9
Likelihood of recommending the IUS at 12 weeks post insertion		
Would recommend	65	89.0
Would not recommend	0	0
Not sure	8	11.0
Alternative method on switching from LNG IUS at 12 months		
No method/ pregnant	5	83.3
Pills	1	16.7

Table 3. Prevalence of side-effects of the IUS at 3- and 12-month post insertion.

Side effects		Prevalence with method at 3 months	Prevalence with method at 12 months
		n (%)	n (%)
Less bleeding	No	90 (91.8)	55 (75.3)
	Yes	8 (8.2)	18 (24.7)
No bleeding	No	92 (93.9)	48 (65.8)
	Yes	6 (6.1)	25 (34.2)
Irregular bleeding	No	66 (67.3)	70 (95.9)
	Yes	32 (32.7)	3 (4.1)
Vaginal bacterial infections	No	96 (98.0)	72 (98.6)
	Yes	2 (2.0)	1 (1.4)
Acne	No	98 (100)	69 (94.5)
	Yes	0 (0.0)	4 (5.5)
Headache/migraine	No	95 (96.9)	100 (100)
	Yes	3 (3.1)	0 (0.0)
Nausea	No	98 (100)	73 (100)
	Yes	0 (0)	0 (0.0)
Pain during sex	No	98 (100)	68 (93.2)
	Yes	0 (0.0)	5 (6.8)
Abdominal discomfort/pain	No	88 (89.8)	69 (94.5)
	Yes	10 (10.2)	4 (5.5)
Breast tenderness/pain	No	97 (99.0)	65 (89.0)
	Yes	1 (1.0)	8 (11.0)
Pelvic discomfort/pain	No	94 (95.9)	73 (100)
	Yes	4 (4.1)	0 (0.0)
Depression	No	98 (100)	69 (94.5)
	Yes	0 (0.0)	4 (5.5)
Mood changes	No	98 (100)	58 (71.2)
	Yes	0 (0.0)	15 (20.5)

*Includes continuers and discontinuers.

presented. Continuation of the use of IUS based on fertility intention was varied among users but higher among users who intended to get pregnant after three years than those intending to become pregnant in under 3 years; this was, also not significant ($p>0.05$). Counseling on side effects was also not statistically associated with continuation of IUS, although

more women who received counseling on side effects (95.5%) than those who did not receive counseling (90%) continued the use of the IUS at 12 months.

Concerning the relationship between satisfaction with method and women's characteristics (Table 6), our result showed that

Figure 1. Percentages for reasons for discontinuation of levonorgestrel intrauterine system (LNG IUS) at 12 months post-insertion.

Table 4. Recommended benefits of the levonorgestrel intrauterine system (LGN IUS) at three months and 12-month post-insertion.

Benefits of IUS recommended	Participant responses at 3 months post-insertion		Participant responses at 12 months post-insertion	
	n	% response	n	% response
Lasts for a long time	43	12.9	-	-
Convenience: once in place nothing else on a regular basis is needed to be done	73	21.9	22	15.1
Discrete: nobody else will know about use	19	5.7	5	3.4
Fewer side effects compared with other methods	79	23.4	21	14.4
Highly effective	78	23.4	48	32.9
Reduced menstrual bleeding	15	4.5	16	11.0
Reversible: possibility of conception in the future	22	6.6	14	9.6
OK for breastfeeding; convenience	2	0.6	6	4.1
Expensive elsewhere and more affordable now	1	0.0	1	0.7
Others	1	0.3	13	8.9

a significantly higher proportion of women, at 12 months, who did not visit a provider because of problems with the method (97.2%) were satisfied with the method compared with those who visited a provider (approximately 76%). This association

also showed statistical significance a 3-months post-insertion. Satisfaction was significantly associated with religious affiliation at 3 months. We found significantly higher satisfaction ($p < 0.05$) with the use of the IUS among users who did not

Table 5. Socio-demographic, family planning and levonorgestrel intrauterine system (LNG IUS) experiences by continuation of the LNG IUS.

User Characteristics	3 months post insertion				12 months post insertion			
	Yes	No	Total	$\chi^2(p)$	Yes	No	Total	$\chi^2(p)$
	n (%)	n (%)			n (%)	n (%)		
Age category (baseline)								
18–24 years old	1 (100.0)	0 (0.0)	1	2.722 (0.436)	1 (100.0)	0 (0.0)	1	3.822 (0.281)
25 – 29 years old	20 (95.2)	1 (4.8)	21		12 (80.0)	3 (20.0)	15	
30 – 34 years old	30 (93.8)	2 (6.2)	32		25 (92.6)	2 (7.4)	27	
35 and older	44 (100.0)	0 (0.0)	44		29 (96.7)	1 (3.3)	30	
Level of schooling (baseline)								
Never attended school	1(100.0)	0 (0.0)	1	1.824 (0.402)	-	-		
Attended but did not complete school					-	-		
Primary	9(90.0)	1 (10.0)	10		7 (100)	0 (0.0)	7	0.693 (0.405)
Secondary or higher	85 (97.7)	2 (2.3)	87		60 (90.9)	6 (9.1)	66	
Marital status (baseline)								
Single	-	-			-	-	-	
Married/living together	93 (96.8)	3 (3.1)	96	0.064 (0.968)	66 (91.7)	6 (8.3)	72	0.091 (0.763)
Widowed	1 (100)	0 (0.0)	1		1 (100)	0 (0.0)	1	
Divorced/separated	1 (100)	0 (0.0)	1					
Religion (baseline)								
Islam	13 (92.9)	1 (7.1)	14	1.211 (0.546)	8 (88.9)	1 (11.1)	9	0.143 (0.931)
Christian (non-Catholic)	69 (97.2)	2 (2.8)	71		49 (92.5)	4 (7.5)	53	
Christian (Catholic)	13 (100)	0 (0.0)	13		10 (100)	0 (0.0)	10	
Parity (baseline)								
0	0	0	0		-	-	-	
1	7 (100)	0 (0.0)	7	0.326 (0.955)	4 (66.7)	2 (33.3)	6	7.224 (0.065)
2	28 (96.6)	0 (3.4)	29		19 (95.0)	1 (5.0)	20	
3	24 (96.0)	1 (4.0)	25		19 (100)	0 (0.0)	19	
4 and more	36 (97.3)	1 (2.7)	37		25 (89.3)	3 (10.7)	28	
Age of youngest child (baseline)								
0 year	34 (97.1)	1 (2.9)	35	0.319 (0.853)	26 (92.9)	2 (7.1)	28	0.659 (0.719)
1–5 years	53 (96.4)	2 (3.6)	55		36 (90.0)	4 (10.0)	41	
6 years and above	8 (100)	0 (0.0)	8		5 (100.0)	0 (0.0)	5	
Pregnancy intention (baseline)								

User Characteristics	3 months post insertion				12 months post insertion			
	Yes	No	Total	$\chi^2(p)$	Yes	No	Total	$\chi^2(p)$
	n (%)	n (%)			n (%)	n (%)		
In less than a year	-	-	-		-	-	-	
In 1–2 years	11 (100)	0 (0.0)	11	2.930 (0.570)	8 (88.9)	1 (11.1)	9	1.973 (0.741)
In 3–5 years	21 (95.5)	1 (4.5)	22		16 (94.1)	1 (5.9)	17	
In more than 5 years	5 (100)	0 (0.0)	5		3 (100.0)	0 (0.0)	3	
Do not know	30 (100)	0 (0.0)	30		22 (95.7)	1 (4.3)	23	
(Never)	28 (93.3)	2 (6.7)	30		18 (85.7)	3 (14.3)	21	
Receiving IUS within 2 weeks of having an abortion or miscarriage								
Yes	2 (100)	0 (0.0)	2	0.064 (0.800)	1 (100.0)	0 (0.0)	1	0.091 (0.763)
No	93 (96.9)	3 (3.1)	96		66 (91.7)	6 (8.3)	72	
Impact of not bleeding to wellbeing[#]								
Positive	-	-	-	-	5 (100.0)	0 (0.0)	5	1.714 (0.424)
Negative	-	-	-	-	5 (71.4)	2 (28.6)	7	
Neutral	-	-	-	-	5 (83.3)	1 (16.7)	6	
Partner feels IUS strings								
Yes	-	-	-	-	17 (94.5)	1 (5.6)	18	0.572 (0.751)
No	-	-	-	-	47 (90.4)	5 (9.6)	52	
Do not know	-	-	-	-	3 (100)	0 (0.0)	3	
Counselling at uptake about what to do if side-effect is experienced								
Yes	33 (97.1)	1 (2.9)	34	0.036 (0.982)	21 (95.5)	1 (4.5)	22	0.693 (0.707)
No	61 (96.8)	2 (3.2)	63		45 (90.0)	5 (10.0)	50	
Do not know	1 (100)	0 (0.0)	1		1 (100)	0 (0.0)	1	
Visits to provider because of problem with method								
Yes	19 (100)	0 (0)	19	0.315 (0.574)	31 (83.8)	6 (16.2)	37	6.361 (0.012)
No	60 (98.4)	1 (1.6)	61		36 (100.0)	0 (0.0)	36	
Total	95 (96.9)	3 (3.1)			67 (91.8)	6 (8.2)		

[#]Among participants who experienced no bleeding.

report vaginal bacterial infection (87.5% satisfied) during the assessment at 12 months. The only case in the sample who reported vaginal infection did not report satisfaction with the method. Not experiencing breast tenderness or pain (90.8%)

and not experiencing acne (88.4%) were also associated with satisfaction with the IUS, compared with experiencing breast tenderness and acne (50% each) with IUS use during the assessment at 12 months post-uptake (Table 7).

Table 6. Socio-demographic, family planning and levonorgestrel intrauterine system (LNG IUS) experiences by satisfaction with the LNG IUS use.

User Characteristics	2 weeks post-insertion (Baseline)			3 months post-insertion*			12 months post-insertion*		
	Not satisfied	Satisfied	Total	Not satisfied	Satisfied	Total	Not satisfied	Satisfied	Total
	n (%)	n (%)	X ² (p)	n (%)	n (%)	X ² (p)	n (%)	n (%)	X ² (p)
Age category (baseline)									
18-24 years old	0(0)	12(100)	12	0(0.0)	1(100)	1	0(0.0)	1(100.0)	1
25 - 29 years old	1(2.4)	41(97.6)	42	4(19.0)	17(81.0)	21	3(18.8)	12(80.0)	15
30 - 34 years old	4(6.5)	58(93.5)	62	8(25.0)	24(75.0)	32	4(15.4)	23(85.2)	27
35 and older	5(5.4)	87(94.6)	92	11(25.0)	33(75.0)	44	3(10)	27(90.0)	30
Level of schooling (baseline)									
Never attended school	0(0.0)	2(100)	2	0(0.0)	1(100)	1	-	-	-
Attended but did not complete school	1(25.0)	3(75.0)	4	-	-	-	-	-	-
Primary	1(6.2)	15(93.8)	16	0(0.0)	10(100)	10	1(14.3)	6(85.7)	7
Secondary or higher	8(4.3)	178(95.7)	186	23(26.4)	64(73.6)	87	9(13.8)	57(86.4)	66
Marital status (baseline)									
Single	0(0.0)	3(100)	3	-	-	-	-	-	-
Married/living together	9(4.5)	189(95.5)	198	23(24.0)	73(76.0)	96	10(13.9)	62(86.1)	72
Widowed	1(33.3)	2(66.7)	3	0(0.0)	1(100.0)	1	0(0.0)	1(100.0)	1
Divorced/separated	0(0.0)	4(100)	4	0(0.0)	1(100.0)	1	-	-	-
Religion (baseline)									
Islam	1(2.7)	36(97.3)	37	2(14.3)	12(85.7)	15	2(20.0)	7(77.8)	9
Christian (non-Catholic)	7(4.6)	144(95.4)	151	21(29.6)	50(70.4)	71	7(13.5)	46(86.5)	53
Christian (Catholic)	2(10)	18(90)	20	0(0.0)	13(100.0)	13	1(10)	10(90)	11

User Characteristics	2 weeks post-insertion (Baseline)				3 months post-insertion*				12 months post-insertion*			
	Not satisfied	Satisfied	Total	$\chi^2(p)$	Not satisfied	Satisfied	Total	$\chi^2(p)$	Not satisfied	Satisfied	Total	$\chi^2(p)$
	n (%)	n (%)			n (%)	n (%)			n (%)	n (%)		
Parity (baseline)												
0	1 (33.3)	2 (66.7)	3	8.488 (0.075)	-	-	-	3.822 (0.281)	-	-	-	3.983 (0.263)
1	2 (12.5)	14 (87.5)	16		1 (16.7)	6 (85.7)	7		2 (40.0)	4 (66.7)	6	
2	3 (5.3)	54 (94.7)	57		7 (24.1)	22 (75.9)	29		4 (19.0)	16 (80)	20	
3	1 (2.1)	47 (97.9)	48		3 (12.0)	22 (88.0)	25		1 (5.3)	18 (94.7)	19	
4 and more	3 (3.6)	81 (96.4)	84		12 (32.4)	25 (67.6)	37		3 (10.7)	25 (89.3)	28	
Age of youngest child (baseline)												
0 year	3 (4)	72 (96.0)	75	0.089 (0.957)	6 (17.6)	29 (82.9)	35	2.284 (0.319)	3 (11.1)	24 (88.9)	27	0.372 (0.830)
1–5 years	5 (4.8)	99 (95.2)	104		16 (29.1)	39 (70.9)	55		6 (15.0)	35 (85.0)	40	
6 years and above	1 (3.8)	25 (96.2)	26		1 (12.5)	7 (87.5)	8		1 (20)	4 (80.0)	5	
Pregnancy intention (baseline)												
In less than a year	0 (0.0)	1 (100.0)	1	10.732 (0.057)	-	-	-		-	-	-	
In 1–2 years	4 (19.0)	17 (81.0)	21		1 (10.0)	10 (90.9)	11	6.659 (0.155)	2 (25.0)	7 (77.8)	9	2.937 (0.568)
In 3–5 years	2 (4.1)	47 (95.9)	49		6 (27.3)	16 (72.7)	22		3 (17.6)	14 (82.4)	17	
In more than 5 years	0 (0.0)	12 (100)	12		0 (0.0)	5 (100)	5		0 (0.0)	3 (100)	3	
Never	2 (3.0)	64 (97.0)	66		5 (16.7)	25 (83.3)	30		4 (17.4)	19 (82.6)	23	
Do not know	2 (3.4)	57 (96.6)	59		11 (36.7)	19 (63.3)	30		1 (4.8)	20 (95.2)	21	
Receiving LNG IUS within 2 weeks of having an abortion or miscarriage (Baseline)												
Yes	2 (25.0)	6 (75.0)	8	7.412 (0.006)	1 (50.0)	1 (50.0)	2	0.800 (0.371)	0 (0.0)	1 (100.0)	1	0.161 (0.688)
No	8 (4.0)	192 (96.0)	200		22 (22.9)	74 (77.1)	96		10 (14.1)	62 (86.1)	72	
Impact of not bleeding to wellbeing*												
Positive	-	-	-	-	-	-	-	-	0 (0.0)	5 (100.0)	5	1.714 (0.424)
Negative	-	-	-	-	-	-	-	-	2 (28.6)	5 (71.4)	7	
Neutral	-	-	-	-	-	-	-	-	1 (16.7)	5 (83.3)	6	

User Characteristics	2 weeks post-insertion (Baseline)				3 months post-insertion*				12 months post-insertion*			
	Not satisfied n (%)	Satisfied n (%)	Total	$\chi^2(p)$	Not satisfied n (%)	Satisfied n (%)	Total	$\chi^2(p)$	Not satisfied n (%)	Satisfied n (%)	Total	$\chi^2(p)$
Partner feels IUS strings												
Yes	-	-	-	-	-	-	-	-	3 (16.7)	15 (83.3)	18	0.613 (0.736)
No	-	-	-	-	-	-	-	-	7 (12.3)	45 (86.5)	52	
Do not know	-	-	-	-	-	-	-	-	0 (0)	3 (100.0)	3	
Counselling at uptake about what to do if side-effect is experienced												
Yes	3 (4.4)	65 (95.6)	68	0.318 (0.853)	5 (14.7)	29 (85.3)	34	2.673 (0.263)	1 (4.5)	21 (95.5)	22	2.500 (0.286)
No	7 (5.1)	128 (94.8)	135		18 (28.6)	45 (71.4)	63		9 (18.4)	41 (81.6)	50	
Do not know	0 (0.0)	5 (100)	5		0 (0.0)	1 (100)	1		0 (0.0)	1 (100.0)	1	
Visits to provider because of problem with method												
Yes	-	-	-	-	3 (17.6)	14 (82.4)	17	7.002 (0.008)	9 (24.3)	28 (75.7)	37	7.165 (0.007)
No	-	-	-	-	1 (1.6)	60 (98.4)	61		1 (2.8)	35 (97.2)	36	
Total (N)	10 (4.8)	198 (95.2)	208		23 (23.5)	75 (76.5)	98		10 (13.7)	63 (86.3)	73	

*Includes continuers and discontinuers.

#Among participants who experienced no bleeding.

Table 7. Satisfaction with levonorgestrel intrauterine system according to experience of side effects at 12-month post-insertion.

Side effects		Satisfaction with method at 12 months*				
		% Not satisfied	% satisfied	Total	X ²	p-value
Less bleeding	No	12.7	87.3	55	0.178	0.673
	Yes	16.7	83.3	18		
No bleeding	No	10.4	89.6	48	1.277	0.258
	Yes	20.0	80.0	25		
Irregular bleeding	No	14.3	85.7	70	0.497	0.481
	Yes	0.0	100.0	3		
Vaginal bacterial infections	No	12.5	87.5	72	6.388	0.011
	Yes	100.0	0.0	1		
Acne	No	11.6	88.4	69	4.717	0.030
	Yes	50.0	50.0	4		
Pain during sex	No	13.2	86.8	68	0.180	0.671
	Yes	20.0	80.0	5		
Abdominal discomfort/pain	No	14.5	85.5	69	0.672	0.412
	Yes	0.0	100	4		
Breast tenderness/pain	No	9.2	90.8	59	10.015	0.002
	Yes	50.0	50.0	4		
Depression	No	14.5	85.5	69	0.672	0.412
	Yes	0.0	100.0	4		
Mood changes	No	17.2	82.5	58	2.997	0.083
	Yes	0.0	100.0	15		

*Includes continuers and discontinuers.

Association analysis between satisfaction with age, the impact of bleeding, pregnancy intention, counseling (Table 6), and side effects like less bleeding, no bleeding, or irregular bleeding (Table 7), were not statistically significant. The findings showed varied high satisfaction with the IUS according to age groupings but were not statistically associated at three months and 12 months post-uptake (Table 6). Satisfaction with the IUS at 12 months post-insertion was markedly higher among women who reported a positive impact of not bleeding from the use of the IUS, compared with those who reported a negative impact of not bleeding, although this was not statistically significant. With regards to pregnancy intention at 12 months post-up-take, women who reported having an intention to get pregnant in the next 3 years and above showed higher satisfaction (range between 82% – 100%) than those who reported having an intention to get pregnant at 2 years (77.8%), this association was not statistically significant.

Experience of satisfaction with the method at 12 months post-uptake was lower among women experiencing less bleeding (83.3%) than women not experiencing less bleeding (87.3%). Similarly, satisfaction was lower with the experience of no bleeding (80%) in contrast to the experience of bleeding (89.6%); these findings on the association between the bleeding changes and satisfaction, were, however, not statistically significant ($p>0.05$).

Discussion

In our study, we found high continuation rates and satisfaction with the hormonal IUD among the users. During the use of contraception, adherence to the method is very important for the prevention of unintended pregnancies, as such the high continuation rate of the IUS contraceptive in our study is indicative of the method's potential and acceptability to reduce the high risk of unwanted and unplanned pregnancies among women

who genuinely need to continue contraception. Similar high continuation rates of the IUS in the first 12 months of use have been reported in previous studies^{12,19}. The IUS has been reported to be highly effective and very tolerable, making the method very acceptable. Side effects are fewer and less serious, which is likely to influence the user's likelihood of satisfaction with and continuation of the method. As our study showed, the majority of the women were satisfied with the IUS, and this satisfaction is correlated to their lowered experience with side effects. Satisfaction from the IUS can also be related to its convenience (in the sense that once the method is inserted, nothing else regularly is needed to be done), as we found that this was among the major reasons the IUS was recommended to other potential FP users.

Experiencing no bleeding or less bleeding, which is a characteristic of the hormonal IUD, presented lower satisfaction (although not statistically significant) with the LNG IUS in our study, compared with other studies where women who experienced amenorrhea tend to appreciate more of this characteristic of the IUS method²⁰. The reason for our findings may not be unconnected to the socio-cultural norms and beliefs that exist in the society about menstruation. Several studies have reported on the misconceptions and myths about menstruation, citing that some women believe menstruation to be a result of the body system discharging unwanted 'bad blood' from the body^{21,22}. This assertion reasonably supports our findings for the lowered satisfaction with the method resulting from reduced or lack of menstrual bleeding. Conversely, we found that most women reporting a negative impact of non-bleeding due to the IUS were less satisfied. Our finding that older women (≥ 35 years) and women with more children (≥ 2 children) presented higher proportion to continue the use of the IUS, compared with their younger counterparts and women with fewer children (≤ 1 child), was not surprising. Several studies have also reported high continuation with LARC among women with more children^{23,24}. Clinicians should encourage younger women and women with fewer children to use LARC as long as they need to delay pregnancies.

Adequate counseling and visiting a health provider for the uptake of health services has been linked with adherence to the use of the health service and of FP specifically^{25,26}. This is important because patients/clients receive relevant information, access check-ups, and get informed about potential risks, fears, and misunderstandings that might exist and can affect the continuation of health services. We found in our study that more women who received counseling continued the use of the IUS than those who did not, this finding, however, was not statistically significant. Although women are free to discontinue the use of the method at any time, health providers offering support on the management of side effects and counseling on expectations of these side effects of the IUS may also have influenced the continuing use of the method that we observed. Previous studies have, however, reported provider bias in the uptake and use of FP methods to influence the choice of FP method^{27,28}. Health providers, however, need to provide

adequate information and counseling to clients to encourage optional use of an FP method to suit the client's FP needs.

A strength of this study is that the data collection approach via phone interview can be thought to have guaranteed less respondent bias on the disclosure of private information about FP use and contraceptives. The limitation of this study, however, is the high attrition rate. Analysis showed some significant difference in the data between the age of participants between baseline and at midline, but not end line of the study, indicating that there was no serious potential attrition bias. Further, while the sampling for this study was widespread and done from 40 facilities across 17 states in Nigeria, given the small sample size and selection of facilities only from the private sector, caution should be taken on the generalization of the findings regarding users of the hormonal IUD in other regions in the country or across the entire country. Further studies are required on the reproductive and health profiles of the users of the IUS in Nigeria with larger sample size and control to improve retention rate. A comparative study with other LARC methods will also provide further insights into the full potential of the IUS method. Evidence is also required on the use of the method in Nigeria for the treatment of menorrhagia and dysmenorrhea, which is quoted as a characteristic of the IUS method, to fully take advantage of these benefits.

Conclusion

In conclusion, the hormonal IUD has positive potential as an acceptable intrauterine contraceptive given the high continuation rate and user satisfaction with the method. The hormonal IUD method presents few side effects which influenced continuation and satisfaction with the method and makes it highly recommended to other potential users of contraceptives. National action is, therefore, recommended for expanding the availability of the hormonal intrauterine system method in the public and private sector to reduce the unmet need for family planning in Nigeria.

Data availability

Underlying data

Figshare: Continuation and user satisfaction of the levonorgestrel intrauterine system (LNG IUS) contraceptive in Nigeria, <https://doi.org/10.6084/m9.figshare.12988313.v11>²⁹.

Extended data

Figshare: Continuation and user satisfaction of the levonorgestrel intrauterine system (LNG IUS) contraceptive in Nigeria <https://doi.org/10.6084/m9.figshare.12988313.v11>²⁹.

This project contains the following extended data:

- Questionnaire and data codes_Continuation_user satisfaction of LNG IUS study

Data are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

Acknowledgments

The authors would like to thank the women who agreed to take part in this study, and the health providers in the SFH Healthy Family Network social franchise who assisted in the recruitment of the women who participated in the study. We

thank Kendal Danna and other members of the Population Service International - Washington D.C, for their expertise and contribution in supporting the implementation of the LNG IUS service delivery and the study in Nigeria with the Society for Family Health. Population Service International was also the co-principal investigator on the study in Nigeria.

References

- National Population Commission (NPC) [Nigeria], ICF: **Nigeria Demographic and Health Survey 2018**. Abuja, Nigeria, and Rockville, Maryland, USA: NPC and ICF, 2019.
[Reference Source](#)
- Thobani R, Jessani S, Azam I, et al.: **Factors associated with the discontinuation of modern methods of contraception in the low income areas of Sukh Initiative Karachi: A community-based case control study**. *PLoS One*. 2019; **14**(7): e0218952.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ali MM, Cleland IH, Shah JG: **Causes and consequences of contraceptive discontinuation: evidence from 60 Demographic and Health Surveys**. *World Heal. Organ*. Geneva, 2012.
[Reference Source](#)
- Rademacher KH, Solomon M, Brett T, et al.: **Expanding access to a new, more affordable levonorgestrel intrauterine system in Kenya: Service delivery costs compared with other contraceptive methods and perspectives of key opinion leaders**. In *Glob Health Sci Pract*. 2016; **4** Suppl 2(Suppl 2): S83–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Riley C, Garfinkel D, Thanel K, et al.: **Getting to FP2020: Harnessing the private sector to increase modern contraceptive access and choice in Ethiopia, Nigeria, and DRC**. *PLoS One*. 2018; **13**(2): e0192522.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Family Planning Evidence Brief – Expanding contraceptive choice. 2018.
[Reference Source](#)
- Contraceptive Discontinuation: Reasons, Challenges, and Solutions**. Family Planning, 2020; [Accessed: 07-Jun-2020].
[Reference Source](#)
- N. B. of S. NBS, U. N. C. F. UNICEF: **Multiple Indicator Cluster Survey 2016-17 Survey Finding Report**. *Mult. Indic. Clust. Surv.* 2016-17, *Surv. Find. Report*. Abuja, Niger. *Natl. Bur. Stat. United Nations Child. Fund*. 2017.
[Reference Source](#)
- Contraceptive Continuation rates**. DataForImpactProject. [Accessed: 19-Nov-2021].
[Reference Source](#)
- Eva G, Nanda G, Rademacher K, et al.: **Experiences with the levonorgestrel intrauterine system among clients, providers, and key opinion leaders: A mixed-methods study in Nigeria**. *Glob Health Sci Pract*. 2018; **6**(4): 680–692.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Suleiman BU, Garba AZ, Makarfi UA, et al.: **The use of levonorgestrel releasing intrauterine system at Ahmadu Bello University Teaching Hospital, Zaria**. *Niger J Basic Clin Sci*. 2017; **14**(1): 30–33.
[Publisher Full Text](#)
- Rademacher KH, Sripathana T, Pfitzer A, et al.: **A global learning agenda for the levonorgestrel intrauterine system (LNG IUS): Addressing challenges and opportunities to increase access**. *Glob Health Sci Pract*. Johns Hopkins University Press, 2018; **6**(4): 635–643.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hubacher D: **The levonorgestrel intrauterine system: Reasons to expand access to the public sector of Africa**. *Glob Health Sci Pract*. 2015; **3**(4): 532–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Beatty MN, Blumenthal PD: **The levonorgestrel-releasing intrauterine system: Safety, efficacy, and patient acceptability**. *Ther Clin Risk Manag*. 2009; **5**(3): 561–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Díaz J, Faúndes A, Díaz M, et al.: **Evaluation of the clinical performance of a levonorgestrel-releasing IUD, up to seven years of use, in Campinas, Brazil**. *Contraception*. 1993; **47**(2): 169–75.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Sivin I, Stern J, Coutinho E, et al.: **Prolonged intrauterine contraception: A seven-year randomized study of the levonorgestrel 20 mcg/day (LNG 20) and the Copper T380 Ag IUDS**. *Contraception*. 1991; **44**(5): 473–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Creinin MD, Jansen R, Starr RM, et al.: **Levonorgestrel release rates over 5 years with the Liletta® 52-mg intrauterine system**. *Contraception*. 2016; **94**(4): 353–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Nanda K, Callahan R, Dorfinger L: **Addressing gaps in the contraceptive method mix: Methods in development**. *Womens Health (Lond)*. 2015; **11**(6): 729–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hubacher D, Masaba R, Manduku CK, et al.: **The levonorgestrel intrauterine system: Cohort study to assess satisfaction in a postpartum population in Kenya**. *Contraception*. 2015; **91**(4): 295–300.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Heikinheimo O, Inki P, Schmelter T, et al.: **Bleeding pattern and user satisfaction in second consecutive levonorgestrel-releasing intrauterine system users: Results of a prospective 5-year study**. *Hum Reprod*. 2014; **29**(6): 1182–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Adinma ED, Adinma JIB: **Perceptions and practices on menstruation amongst Nigerian secondary school girls**. *Afr J Reprod Health*. 2008; **12**(1): 74–83.
[PubMed Abstract](#)
- Rizvi N, Ali TS, et al.: **Misconceptions and Mismanagement of Menstruation among Adolescents Girls who do not attend School in Pakistan**. *J Asian Midwives*. 2016; **3**(1): 46–62.
[Reference Source](#)
- Anguzu R, Sempeera H, Sekandi JN: **High parity predicts use of long-acting reversible contraceptives in the extended postpartum period among women in rural Uganda**. *Contracept Reprod Med*. 2018; **3**: 6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Peipert JF, Zhao Q, Allsworth JE, et al.: **Continuation and satisfaction of reversible contraception**. *Obstet Gynecol*. 2011; **117**(5): 1105–1113.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Shahidzadeh-Mahani A, Omidvari S, Baradaran HR, et al.: **Factors affecting quality of care in family planning clinics: A study from Iran**. *Int J Qual Heal Care*. 2008; **20**(4): 284–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Dehingia N, Dixit A, Averbach S, et al.: **Family planning counseling and its associations with modern contraceptive use, initiation, and continuation in rural Uttar Pradesh, India**. *Reprod Health*. 2019; **16**(1): 178.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sieverding M, Schatzkin E, Shen J, et al.: **Bias in contraceptive provision to young women among private health care providers in south west Nigeria**. *Int Perspect Sex Reprod Health*. 2018; **44**(1): 19–29.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Solo J, Festin M: **Provider bias in family planning services: A review of its meaning and manifestations**. *Glob Health Sci Pract*. 2019; **7**(3): 371–385.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nwala A, Udoh EE, Anyanti J, et al.: **Continuation, Satisfaction LNGIUS Nigeria Dataset (CSV) and Questionnaire with codes**. [Online]. [Accessed: 28-Sep-2020].
https://figshare.com/articles/dataset/_I2988313